

# DISCHARGE INSTRUCTIONS

Ashley E. Ross, M.D., Ph.D.

## **PRESCRIPTIONS**

You will receive prescriptions before discharge. These will include a medication to help manage pain (typically oxycodone). You may want to get that filled before leaving the hospital. Viagra is sometimes given. This can be started approximately one month post operatively and may help with return of erections after surgery. Viagra should not be taken if you are on nitrates for heart disease. If you have heart disease, check with your cardiologist before taking Viagra. Insurance companies generally will not reimburse for more than 6 Viagra per month, so you will have to purchase some of it. Cipro is to be taken twice daily once the catheter is removed to prevent infection. Begin Cipro on the night before you are to have the catheter removed.

## **DIET**

You may eat and drink whatever you wish. Alcohol consumption in moderation is acceptable. Adjust your diet so that you avoid constipation. If you have a problem with constipation you can take Colace, an over the counter stool softener, for prevention after you leave the hospital. If you do become constipated take mineral oil or milk of magnesia. It is important to drink plenty of fluids while the catheter is in place, enough to keep the urine in the tubing (just past the catheter) clear. The urine in the collection bag will almost always be blood tinged, but that is not important as long as the urine in the tubing is pink to clear.

## **AMBULATION**

After you are discharged from the hospital you should avoid heavy lifting (more than 10lbs) and vigorous exercise (calisthenics, golf, tennis, vigorous walking) for total of 4 weeks from the day of surgery. After that you can gradually build up to your pre surgical level of activity but do this gradually. Do not ride a bike for at least 8 weeks from the date of surgery. You may climb stairs slowly. Take frequent short walks during the day (6-8) for 5 minutes or so (like you did in the hospital) while the catheter is in place. After catheter removal, there is no limitation on walking.

While the catheter is in place, you will be more comfortable if you sit in a semi-recumbent position (in a reclining chair, on a sofa, or in a comfortable chair with a footstool). This accomplishes 2 goals: 1) it elevates your legs, thereby improving drainage from veins in your legs, which will reduce the possibility of clot formation (see below); and 2) it avoids placing weight on the area of your surgery in the perineum (the space between the scrotum and the rectum).

There are no other serious restrictions. You may take off the support stockings 1 week after catheter removal, and you may drive your car several days after catheter removal.

## **RETURN TO WORK**

You can work from home when you get home. Most patients who do sedentary “office” activities return to work gradually beginning around 2-4 weeks from the date of surgery. If you do strenuous work (e.g., heavy lifting) then you should wait 4-8 weeks from the date of surgery to return. For those men who travel a lot for business, it is reasonable to wait 4 weeks before returning to a busy travel schedule.

You can drive a car after catheter removal. You will not have your “normal” stamina for up to 3-6 months from the date of surgery, so use common sense in returning to pre surgical activity. Activities that seemed effortless prior to surgery will bring on fatigue more quickly and you may need to rest some during the day.

## **PROBLEMS**

***Bleeding:*** It is not uncommon to have a bloody discharge around the catheter when you strain to have a bowel movement; do not become concerned; it will stop. Also, do not worry about some blood in the urine; it may arise from vigorous walking, or it may occur spontaneously. If this occurs drink plenty of fluids. This will dilute out the blood so that it does not clot off the catheter and will encourage the cessation of bleeding. Blood in the urine usually has no significance and spontaneously resolves on its own.

**IF YOU ARE HAVING BLOOD AROUND OR THROUGH THE CATHETER IN THE 2 DAYS PRECEEDING CATHETER REMOVAL CONTACT ME THROUGH MY OFFICE. THIS MAY BE A SIGN THAT YOUR CATHETER IS NOT READY TO BE REMOVED**

***Leakage around the catheter:*** This is very common, especially when you're up walking around. The tip of the catheter is not in the most dependent part of the bladder; the balloon that holds the catheter in the bladder elevates the tip of the catheter away from the bladder neck. For this reason, when walking many patients have leakage around the catheter. This can usually be managed through the use of diapers or other absorbent materials. If your catheter stops draining completely, lie down flat and drink a lot of water. If after 1 hour there is no urine coming through the catheter tubing, it is possible that your catheter has become obstructed or dislodged. At that point call me (see below).

***Bladder Spasm:*** While the catheter is in place, it is not unusual to have a strong sudden desire to urinate, with pain over the bladder area and simultaneous leakage of urine or blood around the catheter. This is called a bladder spasm and commonly occurs at the time of a bowel movement. If it occurs you should lie down until the discomfort passes. If bladder spasm becomes frequent and bothersome, Motrin or Advil can be used to help stop the spasm. These medicines should not be used if the urine is still bloody because they could lead to clotting of the catheter.

***Wound:*** You may shower after leaving the hospital. The water will not harm the incision or the catheter. You may want to cover up the tape holding the catheter to your leg with a "baggie" or plastic wrap to avoid getting the tape wet.

Some patients develop drainage from the wound once they go home. This can either be clear fluid (a seroma) or a mixture of blood and pus. In either instance it usually can be treated simply. If the wound should open or the edges separates, obtain some hydrogen peroxide and Q-tips; soak the Q-tip in the hydrogen peroxide and place it through the opening in the wound to clean the open area and then remove the Q-tip. This will keep the opening from closing until all the material has drained. I suggest that you shower in the morning washing this area thoroughly (you cannot hurt it). After your shower use the Q-tip and then place a dressing or band-aid over the site. Repeat the Q-tip and dressing before you go to bed that night. Feel free to call me for further advice (see below).

***Clots in the legs:*** During the first 4-6 weeks after surgery, the major complication that occurs in 1-2% of men is a clot in a vein deep in your leg (deep venous thrombosis). This can produce pain in your calf or swelling in your ankle or leg. These clots may break loose and travel to the lung, producing a life threatening condition known as pulmonary embolus.

A pulmonary embolus can occur without any pain or swelling in your leg- the symptoms are chest pain (especially when you take a deep breath), shortness of breath, the sudden onset of weakness or fainting, and/or coughing up blood. If you develop any of these symptoms or pain/swelling in your leg, call me. Also, you should immediately call your local physician or get to an emergency room and state that you need to be

evaluated for deep venous thrombosis or pulmonary embolism. If the diagnosis is made early, treatment with anticoagulation is easy and effective.

If you are traveling home from the hospital in a car, stop the car every 45 minutes and walk around the car to prevent the blood from pooling in the legs. If you are traveling by air, walk the length of the airplane at 45 minute intervals.

**Urinary Tract Infection:** Urinary tract infections can occur with a catheter in place. They can be manifested in several ways. Before the catheter is removed the urine may become permanently cloudy (see below), there may be purulent (thick) drainage around the catheter, and there may be continuous pain at the end of the urethra. This suggests that you may have a urinary tract infection (drainage of mucous around the catheter is normal). Also, it is not unusual for some bacteria to be present in the urine. For this reason, many urologists will place you on an antibiotic for a few days after the catheter has been removed, and I have enclosed a prescription (antibiotic, Cipro) for you to take. Note that it is common to have burning with urination after catheter removal (from irritation of the urethral lining), but this does not mean that you have a urinary tract infection. The burning should improve within several days. It is also common to see passage of some blood or blood clots after catheter removal, but this is of no concern unless it is persistent.

**Urinary sediment:** It is not uncommon to see some sediment in the urine. This can be manifested in a number of different ways. Old clots may appear as dark particles that occur after the urine has been grossly bloody. With hydration these will usually clear spontaneously and are of no concern. Also, the pH (acidity or alkalinity) of the urine changes throughout the day. After a meal the urine oftentimes becomes alkaline. There are normal substances in the urine called phosphates. They precipitate out in alkaline urine and form cloudy masses in the urine. If you see these periodically do not be concerned. This is a normal phenomenon. Finally, if the urine is persistently cloudy this suggests that an infection may be present (see above).

**Pain:** Abdominal pain is common. The pain is from irritation of the abdominal muscles; sometimes it is where the drainage tube exited. It will resolve spontaneously, but it is not uncommon to have sensitivity around the incisions for 3-6 months after surgery. **You may take the pain pills that I prescribed for you, but I encourage patients to start with ibuprofen (if you do not have trouble with ulcers) or Tylenol. I usually recommend taking 3 200mg tablets of ibuprofen (Advil or Motrin) three times a day with meals for the first 5 or 6 days. This provides excellent pain control with minimal side effects. After 5 or so days, wean off the ibuprofen as this can be harsh on the GI tract. If you take Prevacid/Protonix/Nexium, etc. be sure you resume this after surgery.** You may notice sensitivity when you fasten your pants belt or a seat belt. This is normal. Finally, you may notice firm areas or lumps in the incision. This is part of the normal healing process. If you notice a hard area or lump at the top of the incision (near the umbilicus), this is where the suture material was tied and is also normal. It will resolve with time.

It is very common to have a deep feeling of discomfort in the perineum (between the scrotum and rectum), especially after sitting. The pain is coming from the area where the operation took place and will disappear with time but may be present for 1-2 months after surgery. Avoid sitting for a long time if it is bothersome or sit on a “doughnut” (round cushion). Discomfort in the testicles is very common after radical prostatectomy because the spermatic cord (attached to the testicle) is stretched during the operation. This discomfort will disappear in time but can last 3-6 months after surgery. If bothersome, use Motrin or Advil if the urine is clear.

**Swelling:** It is very common to have swelling and discoloration of the scrotum and the penile skin after radical prostatectomy. This is simply fluid that has not been absorbed by the body. It is not harmful. If the scrotum is swollen, put a rolled hand towel underneath the scrotum to elevate it when lying down.

***Catheter removal:*** Your catheter should be removed approximately 10 to 14 days from the day of surgery. On the day you are going to have your catheter removed drink a lot of fluids before you arrive at the office. On that day I am only concerned with whether or not you are urinating with a strong stream. The recovery of urinary control takes longer (see below). It is very important for me to write to the physician that is removing your catheter (if you are not returning to me for removal). If you cannot give me the name and address while you are in the hospital, please call my secretary at 469-626-5704 and give her the information so this communication takes place before you arrive at the office of the physician removing the catheter. **IF YOU ARE HAVING BLOOD AROUND OR THROUGH THE CATHETER IN THE 2 DAYS PRECEEDING CATHETER REMOVAL CONTACT ME THROUGH MY OFFICE. THIS MAY BE A SIGN THAT YOUR CATHETER IS NOT READY TO BE REMOVED.**

### **COMMUNICATION WITH DR. ROSS – SEE CHART ON LAST PAGE FOR OVERVIEW**

Call my office after you are discharged from the hospital to set up or confirm our first appointment. We will have frequent contacts during the first year after surgery (typically 3 month intervals for 1<sup>st</sup> year after surgery). Many of these can be through telephone appointments that will be scheduled for 5-10 minutes

**If you have any problems when you are at home, call me at the office between 8:30 a.m. and 4:30 p.m. My secretary will arrange for us to speak. If it is an emergency my secretary will get in touch with me. If you have an emergency at night or on a weekend call me at 410-782-5068.**

### **URINARY CONTROL**

Problems with urinary control are common once the catheter is removed. Do not become discouraged. Urinary control returns in 3 phases: Phase I – you are dry when lying down at night; Phase II – you are dry when walking around; Phase III – you are dry when you rise from a seated position. This is the last component of continence that returns. In the early phases your urinary stream may be weak if the bladder is not filling and most of the urine is leaking into a pad. You may also experience more frequent urination after surgery, as the bladder capacity increases over time. Everyone is different and, for this reason, I cannot predict when you will be dry. To speed up your recovery, practice stopping and starting your urinary stream every time you void. To do this, you must stand up to urinate. To shut off your urinary stream, contract the muscles that you use to keep from passing gas. Until your control returns completely, wear a pad or disposable diaper. You can obtain Depends, and adult diaper, or security pads from your local grocery store or pharmacy.

As urinary control returns, it is not uncommon for patients to continue to wear protective pads for “security” even when they don’t need them. To make sure that you do not become pad dependent unnecessarily, experiment with not using a pad when you are at home and not working. Many patients will have the sensation that they are leaking urine when, in fact, they will find that there has been no leakage on the underwear.

Do not wear an incontinence device with an attached bag, a condom catheter, or a clamp. If you do, you will not develop the muscular control necessary for continence. Until your urinary control returns, avoid drinking excessive amounts of fluids. Also, limit your intake of alcohol and caffeine-both will make the problem worse. Once the catheter is removed, limit fluids to the amount necessary to satisfy your thirst.

It is common for the urinary stream to be slower after surgery. But if you notice a progressive decrease in the force of the urinary stream this could indicate a scar. Do not wait until the urine stream is so slow that you have to strain or push to urinate. Call me if you notice a progressive slowing of the stream since simple dilation of the scar where the bladder and urethra were joined can alleviate the problem if caught early.

If you develop a red painful rash while urinary control is returning, you may have a fungal infection, especially if you were treated with antibiotics. This usually responds well to treatment with Lotrimin cream, a non-prescription formulation that can be purchased over the counter at a pharmacy.

## **SEXUAL FUNCTION**

Erections return gradually (much slower than urinary control) and continue to improve even up to 4 years after surgery. Be patient. As I told you before the operation, the return of sexual function varies depending upon the age of the patient, the extent of the tumor (whether nerves had to be removed), and the level of sexual functioning before the operation. Men who had declining sexual function prior to surgery will have a greater chance of problems with erections after surgery. There are some patients who don't recover potency until two years after surgery. Erections return gradually and quality improves month by month with effort.

After surgery, it is important for men to have realistic expectations of the quality of erections. At first erections will be partial and not likely strong enough for penetration. But a partial erection is success! Open a bottle of champagne if you get a partial erection because with continued effort they will get strong enough for penetration. Most men do not have recovery of an erection that is "exactly" the same as before surgery. Men who recover erections strong enough for intercourse usually have erections that are more difficult to attain and maintain, and because of this it is common for libido (desire for sexual activity) to decrease.

The stimuli for erection during the first year will be different. Visual and psychogenic stimuli will be less effective and tactile sensation will be more effective. Indeed, the major stimulus for erections during the first year postoperatively is tactile sensation. For this reason, do not be afraid to experiment with sexual activity - you can do no harm. If you obtain a partial erection, attempt vaginal penetration; many patients find that erections are maintained better when upright (rather than lying down) and that vaginal penetration is easier from behind. Lubrication of the vagina with K-Y jelly can help. Vaginal stimulation will be the major factor that encourages further erections. Do not wait until you have the "perfect erection" before attempting intercourse. In addition, you should be able to have an orgasm even if you do not have an erection. With orgasm there will be no emission of semen because the prostate and seminal vesicles have been removed.

When erectile function begins to return many patients complain that they lose their erections when they attempt intercourse. This is caused by a venous leak. This can be overcome by placing a soft tourniquet at the base of the penis before foreplay. The purpose of this tourniquet is to retain the blood in the penis once blood flow increases secondary to stimulation. Do not worry; the tourniquet will not impede the flow of blood into the penis. My patients have told me that rubber bands, ponytail holders, or "erection rings" (which can be obtained from novelty stores) work. The best product is made by Vivus and is called the Actis venous flow controller. You can read about it on the web site [Vivus.com](http://Vivus.com) under product information and you can order by calling the number listed on the web site.

Viagra (or another PDE5 inhibitor like Cialis or Levitra) can be a very effective aid to improve erections during the recovery period. Do not take this medication if you are on nitrate to treat heart disease (coronary artery disease). Once you are ready to begin sexual activity, I suggest that you take a 100mg tablet 1-2 hours prior to activity on an empty stomach. Do not use Viagra more than once daily.

It is reasonable to begin experimenting with sexual activity after catheter removal whenever you feel ready. Do not wait for erections to return on their own-they will not without a lot of persistence and perseverance on the part of both partners. Patients who are willing to continue attempts to produce erections- despite lack of a perfect erection- are more likely to have return of erectile function in the long run. Begin experimenting with erections as soon as possible after catheter removal and this will increase the likelihood for recovery in the long-term (use it or lose it). If you are not having success with return of erections, we will talk about other options for management during our telephone appointments.

**LONG-TERM EVALUATION – SEE LAST PAGE FOR OVERVIEW OF FOLLOWUP**

I would like to speak to you by phone 2 weeks after catheter removal, then 3 months, 6 months, 9 months and 1 year from the date of surgery. You should have a PSA measurement done 1-3 months from the date of surgery (mail or FAX to me) so we can discuss the result at our 3-month from surgery date conversation. PSA measurement is the only follow-up exam that is needed and should be done at 6-month intervals for the first year after surgery.

**OVERVIEW OF FOLLOWUP**

<b>Time after Surgery</b>	<b>1 month</b>	<b>3 months</b>	<b>6months</b>	<b>12 months</b>	<b>Annually</b>	<b>Anytime</b>
Update on Urinary Control	X	X	X	X	X	X
Update on sexual function		X	X	X	X	X
PSA blood test		X	X	X	X	X

PSA blood tests should be obtained 2 weeks prior to scheduled phone call or office visit.